

$\mathcal{W}$ GEOMETRY FROM FEDOSOV'S DEFORMATION QUANTIZATION

Carlos Castro

Center for Theoretical Studies of Physical Systems,
Clark Atlanta University, Atlanta, Georgia 30314

January, 1998

ABSTRACT

A geometric derivation of W_∞ Gravity based on Fedosov's deformation quantization of symplectic manifolds is presented. To lowest order in Planck's constant it agrees with Hull's geometric formulation of classical nonchiral W_∞ Gravity. The fundamental object is a $\mathcal{W}$ -valued connection one form belonging to the exterior algebra of the Weyl algebra bundle associated with the symplectic manifold. The $\mathcal{W}$ -valued analogs of the Self Dual Yang Mills equations, obtained from a zero curvature condition, naturally lead to the Moyal Plebanski equations, furnishing Moyal deformations of self dual gravitational backgrounds associated with the complexified cotangent space of a two dimensional Riemann surface. Deformation quantization of W_∞ Gravity is retrieved upon the inclusion of all the $\hbar$ terms appearing in the Moyal bracket. Brief comments on Non Commutative Geometry and M(atrrix)theory are made.

Keywords : Integrable systems, star products, $\mathcal{W}$ -geometry, Moyal-Fedosov quantization, strings.

Subj. Class : Differential Geometry

1991 MSC : 32 C34, 34 C35, 58 H15.

I. Introduction

Since the classic work of Bayen et al [1] Moyal deformation quantization techniques [2] are starting to become very relevant in the area of Non Commutative Geometry, for example (see [14] for a current review). W_∞ algebras, strings, gravity, etc were very popular candidates to extensions of the ordinary two-dimensional Conformal Field Theory (CFT) description of strings based on Kac-Moody and Virasoro algebras . For an extensive review on higher conformal spin extensions of CFT we refer to the Physics Reports article [5]. Not long ago, the author [8] was able to show that non-critical W_∞ strings are devoid of BRST anomalies for target spacetimes of dimension $D = 27$. The supersymmetric case yielded $D = 11$ and we suggested that the an anomaly free (super) membrane should support non-critical W_∞ strings in their spectrum.

W_∞ covariance is of crucial importance in the Weyl-Wigner-Groenewold-Moyal quantization process [4] and for this purpose its relevance should be investigated further. The role of W_∞ algebras in Moyal quantization was also investigated by [17]. The authors in [24] were able to realize that nonlinear W_∞ algebras could be obtained by Moyal deformations of the classical w_∞ linear ones. Other important connections among membranes and strings were raised in [25].

In this letter we present a very natural framework, in the language of Fedosov's deformation quantization [3], where the original formulation of W_∞ Geometry presented by Hull [6] can be incorporated in a very simple fashion. In section **II** we discuss Fedosov's Geometry and Moyal SDYM theories in R^4 . In **III** we review Hull's work and show how his formulation admits a straightforward incorporation in the language of Moyal-Fedosov quantization. In the final concluding section we outline some of the many applications that deformation quantization techniques have in the theory of extended objects.

II. Fedosov Geometry and Moyal Self Dual Yang Mills

2.1. Fedosov's Deformation Quantization

In essence, Fedosov's [3,33] deformation quantization ; i.e deformations of the Poisson Lie algebra structure on a symplectic manifold, is a generalization of the Bayen et al [1] and the Weyl, Wigner, Groenewold, Moyal (WWGM)[2] deformation quantization on flat phase spaces. Given a symplectic manifold, $\mathcal{M}$ of dimension $2n$, with a symplectic globally defined non-degenerate two-form ω , allows to define a symplectic structure in each tangent space, $T_x\mathcal{M}$ to the point x . The Weyl algebra, $\mathcal{W}_x$ corresponding to the symplectic space, $T_x\mathcal{M}$, is the associative non-commutative algebra over $\mathbf{C}$ with a unit element. The elements of the Weyl algebra are defined by a formal power series :

$$a(y) = \sum h^k a_{k,i_1,\dots,i_l} y^{i_1} \dots y^{i_l}. \quad 2k + l \geq 0. \quad (1)$$

where h is a formal parameter which can be identified with $\hbar$ and the coordinates $y^1, \dots, y^{2n} \in T_x\mathcal{M}$ are associated to a tangent vector at the point x . The degrees 1, 2 are prescribed for the variables y, h respectively, so one is summing in eq-(1) over elements of the algebra of non-negative degree. The noncommutative product on the Weyl algebra, $\mathcal{W}_x$, which determines its associative character, of two elements, $a(y), b(y)$ is defined :

$$a(y) \circ b(y) = \sum_{k=0}^{\infty} \left(\frac{i\hbar}{2}\right)^k \frac{1}{k!} \omega^{i_1 j_1} \dots \omega^{i_k j_k} (\partial_{i_1} \dots \partial_{i_k} a) (\partial_{j_1} \dots \partial_{j_k} b). \quad (2)$$

where ω^{ij} are the components of the inverse tensor of ω_{ij} , the symplectic two-form.

Having defined the Weyl algebra at a point $x \in \mathcal{M}$ one can define an algebra bundle structure by taking disjoint unions of the Weyl algebras at each point; i.e a fibrewise addition. In this fashion a Weyl algebra bundle, $\mathcal{W}$, is constructed as x runs over all the points in $\mathcal{M}$. This allows to build in a set of sections, $\mathcal{E}(\mathcal{W})$, denoted by $a(x, y, \hbar)$ which can be written in a power series :

$$a(x, y, \hbar) = \sum h^k a_{k,i_1,\dots,i_l}(x) y^{i_1} \dots y^{i_l}. \quad 2k + l \geq 0. \quad (3)$$

where now $a_{k,i_1,\dots,i_l}(x)$ are a set of smooth functions defined on the symplectic manifold. Differential forms can be constructed whose elements are $\mathcal{W}$ -valued (instead of the customary forms taking values in a Lie algebra). A $\mathcal{W}$ -valued q differential form may be written :

$$A(x, y, \hbar) = \sum h^k a_{k; i_1, \dots, i_l; j_1, \dots, j_q}(x) y^{i_1} \dots y^{i_l} dx^{j_1} \wedge \dots \wedge dx^{j_q}. \quad 2k + l \geq 0. \quad (4)$$

and an exterior algebra extension of the Weyl algebra bundle, $\mathcal{E}(\mathcal{W} \otimes \Lambda)$ is obtained by constructing a deformation of the wedge product, $\wedge^\circ$. Differential operators analogous to the exterior derivative and its dual (divergence) are defined in [3] and reviewed in [10,33]. We shall not repeat it here.

A torsion-free symplectic connection, ∂ , preserving the symplectic structure can also be defined. The symplectic connection does *not* change the degree of the Weyl algebra bundle. $\partial : \mathcal{E}(\mathcal{W}_p \otimes \Lambda^q) \rightarrow \mathcal{E}(\mathcal{W}_p \otimes \Lambda^{q+1})$. Fedosov defined a more general connection, D , the analog of a gauge covariant derivative, acting in an element, a , of the Weyl algebra bundle as :

$$Da = \partial a + \frac{1}{i\hbar}[\gamma, a]. \quad \gamma \in \mathcal{E}(\mathcal{W} \otimes \Lambda^1). \quad (5)$$

for γ a globally defined $\mathcal{W}$ -valued one form. Since every symplectic manifold can be equipped with an almost complex structure, J , such that the tensor defined by $g(X, Y) = -\omega(JX, Y)$ for all vector fields, X, Y , is a Riemannian metric on $\mathcal{M}$. This yields the curvature R associated with the symplectic connection : $R = \frac{1}{4}R_{ijkl}y^i y^j dx^k \wedge dx^l$. The commutator is defined as : $[a, b] = a \wedge^\circ b - (-1)^{qp} b \wedge^\circ a$ where a, b are respectively $\mathcal{W}$ valued q, p differential forms. The curvature of the connection, D , satisfies the property :

$$D^2 a = \frac{1}{i\hbar}[\Omega, a]. \quad \Omega = R + \partial\gamma + \frac{1}{i\hbar}\gamma^2. \quad (6)$$

A connection D is Abelian if, and only if, for any section, $a \in \mathcal{E}(\mathcal{W} \otimes \Lambda^1)$, $D^2 a = 0$. Hence from (6) one can infer that the curvature of every abelian connection, Ω , is central (commutes with all the elements a of the Weyl algebra) and is proportional to $\omega_{ij} dx^i \wedge dx^j$. Abelian connections are very relevant in the construction of the algebra of Quantum Observables, $\mathcal{E}(\mathcal{W}_D)$, which is the subalgebra of the Weyl algebra bundle comprised of *flat sections*,

$$a \in \mathcal{E}(\mathcal{W}_D) \Rightarrow D^2 a = 0. \quad (7)$$

these flat sections (with respect the Abelian connection) generate a subalgebra of the Weyl algebra bundle called the algebra of Quantum Observables. It is the analogus of BRST invariant states in string theory.

Finally one may establish the relationship with the ordinary Moyal star product. Fedosov showed that one can assign to a flat section (relative to an Abelian connection) $a(x, y, \hbar)$ a central element of the Weyl algebra, $a_o(x, y = 0, \hbar) \in \mathcal{Z}$, and viceversa. The bijection map (which we shall not go into details) and the star product in the central elements of the algebra, $a_o, b_o \in \mathcal{Z}$ are :

$$\sigma : \mathcal{W}_D \rightarrow \mathcal{Z}. \quad \sigma(a) = a_o. \quad \sigma^{-1} a_o = a. \quad a_o * b_o = \sigma(\sigma^{-1}(a_o) \circ \sigma^{-1}(b_o)) = \sigma(a \circ b). \quad (8)$$

in this way one can construct a star product in the space of central sections, $\mathcal{Z}$, inherited from the noncommutative associative Fedosov fibrewise product of flat sections belonging to the subalgebra of Quantum Observables. When the symplectic manifold is flat, R^{2n} , the star product reduces then to the ordinary Moyal star product as expected. In this particular case the tangent bundle is trivial, $T(R^{2n}) \sim R^{2n} \times R^{2n}$ and there is no difference between the fibrewise (y^i) Fedosov products and the base space (x^n) Moyal ones. For definitions of the trace of Quantum Observables in the Weyl algebra bundle, its inner automorphisms (analogous to gauge transformation), symplectic diffs,... we refer to the literature [3,10].

2.2 Moyal Self Dual Yang Mills

It has been known for some time to the experts that 4D Self Dual Gravity can be obtained from $SU(\infty)$ SDYM, an effective 6D theory, by dimensional reduction. We refer to [9] for an extensive list of references. The bosonic and supersymmetric case was studied also by the author [7]. Moyal deformations of Self Dual Gravity were proposed by Strachan [18] and rotational Killing symmetry reductions furnished the Moyal quantization of the continuous Toda field [7]. Generalized Moyal Nahm and continuous Moyal Toda equations were developed by [27] based on the work of the SDYM equations by Ivanova and Popov [41]. W_∞ is a natural symmetry of these integrable theories. Takasaki [19] and Strachan, among many others, have emphasized the importance of higher dimensional integrable theories.

In this section we will write down the main equations of Moyal deformations of SDYM theories in R^4 leaving all the technical details for the reference [9,10]. The basic equations are obtained from a zero curvature condition which allows to gauge two of the fields $A_x, A_y = 0$ and yields for the remaining third equation :

$$\partial_{\tilde{x}} A_{\tilde{y}} - \partial_{\tilde{y}} A_{\tilde{x}} + [A_{\tilde{y}}, A_{\tilde{x}}] = 0. \quad (9)$$

A WWGM quantization of a $SU(N)$ SDYM requires finding a representation of the Lie algebra $su(N)$ in a suitable Hilbert space, where the $\hat{A}_\mu$ are vector-valued operators living in a Hilbert space. A WWGM quantization requires the construction of operators acting in the Hilbert space of square integrable functions on the line, $L^2(R)$, and the use of the symbol WWGM map to define a one-to-one correspondence of self adjoint operators into real valued smooth functions in the phase space q, p associated with the line. Evenfurther, the WWGM map takes commutators, $\frac{1}{i\hbar}[\hat{A}, \hat{B}]$ into Moyal Brackets. In [27] we have shown that in general one should enlarge the phase space with the introduction of q^i, p^i to accomodate for the Moyal deformations of continuum Lie algebras introduced by Saveliev and Vershik [42]. Hence eq-(9) becomes an effective 6D equation after the WWGM quantization :

$$\partial_{\tilde{x}} A_{\tilde{y}} - \partial_{\tilde{y}} A_{\tilde{x}} + \{A_{\tilde{y}}, A_{\tilde{x}}\}_{Moyal} = 0. \quad (10)$$

where now, $A_\mu(x, y, \tilde{x}, \tilde{y}, q, p, \hbar)$. The last equation admits, in the $\hbar \rightarrow 0$ limit, reductions to many of the known integrable equations, like the Plebanski first and second heavenly equations, the Park-Hussain and Grant equations,...[9]. Furthermore, eq-(10) may be

obtained directly from a Lagrangian [9]. In fact, dynamics of higher spin fields can be encoded in zero curvature constraints. This has been explained in particular by Vasiliev [40]. The $6D$ Moyal SDYM equations (10) admit a reduction to $4D$ as follows :

$$A_{\tilde{y}} = \partial_x \Theta = \partial_x \Theta' - \frac{1}{2} \tilde{x}. \quad A_{\tilde{x}} = -\partial_y \Theta = -\partial_y \Theta' + \frac{1}{2} \tilde{y}. \quad (11)$$

with the $6D$ function Θ :

$$\Theta(x, y, \tilde{x}, \tilde{y}, q, p, \hbar) \equiv \Theta'(x + \tilde{y}, \tilde{x} - y, q, p, \hbar) - \frac{1}{2}(x\tilde{x} + y\tilde{y}). \quad (12)$$

Inserting (11,12) into (10) yields the $4D$ Moyal first heavenly Plebanski equation :

$$\{\Omega_w, \Omega_{\tilde{w}}\}_{Moyal} = 1. \quad \Omega(w, \tilde{w}, q, p, \hbar) \equiv \Theta'. \quad w = x + \tilde{y}; \quad \tilde{w} = \tilde{x} - y \quad (13)$$

Eq-(13) can also be rewritten, due to the fact that the Moyal bracket of the variables $x, y, \tilde{x}, \tilde{y}$ is zero, :

$$\{\partial_x \Theta(x, y, \tilde{x}, \tilde{y}, q, p, \hbar), \partial_y \Theta(x, y, \tilde{x}, \tilde{y}, q, p, \hbar)\} = 1. \quad (14)$$

The dimensional reduction of the Moyal SDYM theory, from $6D \rightarrow 4D$, leading to the Moyal Plebanski equation, can be interpreted as a foliation of the $6D$ space into $4D$ leaves endowed with Moyal deformed (self dual/anti self dual curvatures) hyper-Kahler Ricci flat metrics and parametrized by the coordinates $\tilde{x}, \tilde{y}$ of the $6D$ space :

$$\{\partial_x K_{\tilde{x}, \tilde{y}}(x, y, q, p, \hbar), \partial_y K_{\tilde{x}, \tilde{y}}(x, y, q, p, \hbar)\}_{Moyal} = 1. \quad (15)$$

where $K_{\tilde{x}, \tilde{y}}(x, y, q, p, \hbar)$ is a two-parameter family of Moyal deformed Kahler potentials. This is attained by setting in (14,15) :

$$\partial_y K_{\tilde{x}, \tilde{y}}(x, y, q, p, \hbar) = \partial_y \Theta(\tilde{x}, \tilde{y} | x, y, q, p, \hbar). \quad \partial_x K_{\tilde{x}, \tilde{y}}(x, y, q, p, \hbar) = \partial_x \Theta(\tilde{x}, \tilde{y} | x, y, q, p, \hbar). \quad (16)$$

Hence, for running values of $\tilde{x} = \tilde{x}_o; \tilde{y} = \tilde{y}_o$ which characterize the foliation, eqs-(13,14,15) yield a two-parameter family of Moyal heavenly metrics encoded in the Moyal Plebanski potential : $\Omega(w, \tilde{w}, q, p, \hbar)$. The foliation is represented by the two-parameter family of four-dimensional noncommutative manifolds, $X_{\tilde{x}, \tilde{y}}^4(x, y, q, p, \hbar)$, since the star product of two functions in phase space is noncommutative. The connections between Non Commutative Geometry, Matrix models [11,12,14,15] and String theory [13] is now being developed by a large number of authors. We apologize for excluding many relevant references. The relevance of Self Dual Gravity in connection with $N = 2$ strings was initiated by [20] and pursued by many others [30,31]. Important remarks about M (atrix) models and $N = 2$ strings have appeared in [29]. The role of self dual gravity and W_∞ algebras was initially worked out, among others, by [21,22,23]. This completes this short review about Moyal SDYM in R^4 .

III. W-Geometry From Fedosov

3.1 Hull's Formulation of W_∞ Gravity

Sometime ago, Hull [6] with great insight, presented a geometric formulation of W_∞ Geometry as a gauge theory of the group of symplectic diffeomorphisms of the cotangent bundle of a two-dimensional Riemann surface, $Diff_o(T^*\mathcal{N})$. The infinite set of symmetric gauge tensor fields are $\tilde{h}_{(n)}^{\mu_1 \dots \mu_n}$, $n = 2, 3, \dots \infty$; $\mu = 0, 1$. These transform as densities under $\mathcal{W}$ gauge transformations :

$$\begin{aligned} \delta \tilde{h}_{(s)}^{\mu_1 \dots \mu_s} = & \sum_{m,n} \delta_{m+n,s+2} [(m-1) \lambda_{(m)}^{(\mu_1 \mu_2 \dots} \partial_\nu \tilde{h}_{(n)}^{\dots \mu_s) \nu} - (n-1) \tilde{h}_{(n)}^{\nu(\mu_1 \mu_2 \dots} \partial_\nu \lambda_{(m)}^{\dots \mu_s)} \\ & + \frac{(m-1)(n-1)}{p-1} \partial_\nu \{ \lambda_{(m)}^{\nu(\mu_1 \mu_2 \dots} \tilde{h}_{(n)}^{\dots \mu_s)} - \tilde{h}_{(n)}^{\nu(\mu_1 \mu_2 \dots} \lambda_{(m)}^{\dots \mu_s)} \}]. \end{aligned} \quad (17)$$

s labels the conformal spin $2, 3, 4, \dots \infty$ of the gauge fields whose physical components have helicity $\pm s$. From the generalized notion of a scalar line element

$$ds = (g_{\mu_1 \dots \mu_n} dx^{\mu_1} \dots dx^{\mu_n})^{(1/n)}$$

Hull proposed an action of the form :

$$S = \int d^2x \tilde{F}(x, y). \quad \tilde{F}(x, y) = \sum_{n=2}^{\infty} \frac{1}{n} \tilde{h}_{(n)}^{\mu_1 \dots \mu_n} y_{\mu_1} \dots y_{\mu_n}. \quad (18)$$

where the function $\tilde{F}(x, y)$ is a co-metric $\mathcal{W}$ -density in $d = 2$ instead of a $\mathcal{W}$ -scalar. The action represents the integrated generalized world interval along a section of the bundle $T^*\mathcal{N}$, where the fiber coordinates, y_μ , when restricted to a section, Σ , can be interpreted as the gradients of the matter fields $y_\mu|_\Sigma = y_\mu(x) = \partial_\mu \phi(x)$. Hull used one and only one bosonic scalar field, $\phi(x)$, living in the two-dimensional world sheet. Later we will see how to include a set of matter fields, $\phi^i(x)$ representing the embedding spacetime coordinates of the string worldsheet. The nonlinear transformation property of the field $\phi(x^\mu)$ is :

$$\delta \phi = \Lambda(x^\mu, y_\mu) = \sum_{n=2}^{\infty} \lambda_{(n)}^{\mu_1 \dots \mu_n}(x^\mu) y_{\mu_1} \dots y_{\mu_n}. \quad (19)$$

However, the formulation based on the infinite number of fields $\tilde{h}_{(s)}^{\mu_1 \dots \mu_s}$ was redundant (the action was reducible) in the sense that there are more gauge fields than are needed. For example, $\tilde{h}^{\mu\nu}$ has three independent components for only two gauge symmetries. Hull proved that a gauge invariant (invariant under the transformations given by (17,19)) constraint could be consistently imposed on all the gauge fields in such a way that one could recast the action solely in terms of a set of unconstrained fields, $h_{(s)}^{\mu_1 \dots \mu_s}$ with their traces removed at the linearized level after exploiting the $\mathcal{W}$ -Weyl conformal invariance. Thus, at the *linearised* level with respect to a flat $\eta_{\mu\nu}$ two-dim metric and to lowest order in the gauge fields one has :

$$\tilde{h}_{(s)}^{\mu_1 \dots \mu_s} = [h_{(s)}^{\mu_1 \dots \mu_s} - traces] + O(h^2) \dots$$

$$\delta \tilde{h}_{(s)}^{\mu_1 \dots \mu_s} = \partial^{(\mu_1} \lambda_{(s)}^{\mu_2 \dots \mu_s)} + \dots; \quad \lambda_{(s)}^{\mu_1 \dots \mu_{s-1}} = [k_{(s)}^{\mu_1 \dots \mu_{s-1}} - traces] + O(k^2) \dots \quad (20)$$

The gauge-invariant background independent master constraint that generates *all* the constraints on the gauge fields, upon expansion in powers of y_μ , and which allows to recast the action solely in terms of the unconstrained fields is :

$$\det[\tilde{G}^{\mu\nu}(x^\mu, y_\mu)] = 1. \quad \tilde{G}^{\mu\nu}(x^\mu, y_\mu) = \frac{\partial^2 \tilde{F}(x^\mu, y_\mu)}{\partial y_\mu \partial y_\nu}. \quad (21)$$

For $2 + 2$ signature one has -1 for the determinant instead in the r.h.s of (21). The constraint (21) can be solved in a particular way by recurring to the $\mathcal{W}$ -Weyl conformal invariance which allows to gauge away all the traces of the unconstrained fields, leaving only traceless fields in the action with helicities $\pm s$.

The geometrical significance of the constraint (21) is the following . If one sets $y_\mu = z_\mu + \bar{z}_\mu$ where z_μ are the complex coordinates on R^4 , $\mu = 0, 1$, allows to view $(x^\mu, z_\mu, \bar{z}_\mu)$ as the coordinates for the bundle $\mathbf{C} T_x^* \mathcal{N}$, the complexification of the cotangent space $T_x^* \mathcal{N}$ at a point $x \in \mathcal{N}$. The co-metric function is then reinterpreted as :

$$\tilde{F}(x^\mu, y_\mu) = K_x(z, \bar{z}) = \tilde{F}(x^\mu, z_\mu + \bar{z}_\mu). \quad (22)$$

with $K_x(z, \bar{z})$ the Kahler potential depending on the combination $z_\mu + \bar{z}_\mu$ which is tantamount of a Killing symmetry reduction condition, the metric does not depend on the two imaginary components of z_μ . :

$$\frac{\partial^2 K_x(z, \bar{z})}{\partial z_\mu \partial \bar{z}_\nu} = \frac{\partial^2 \tilde{F}(x^\mu, y_\mu)}{\partial y_\mu \partial y_\nu}. \quad (23)$$

The gauge invariant (under the transformations given by (17,19)) master constraint (21) was equivalent to the Monge-Ampere equation (Plebanski equation) for a Ricci flat, hyper-Kahler manifold associated with the complexified cotangent space at each point x^μ , of the original two-dimensional surface, world-sheet, $\mathcal{N} : \mathbf{C} T_x^* \mathcal{N} \sim \mathbf{C}^2$. This implies that for each x^μ , the corresponding curvature tensor is either self-dual or antiselfdual. Summarizing, for each, $x^\mu \in \mathcal{N}$, $\tilde{F}(x^\mu, y_\mu = z_\mu + \bar{z}_\mu) = K_x(z, \bar{z})$ is the Kahler potential for a hyperKahler metric on R^4 with two commuting tri-holomorphic Killing vectors. Hence, $K_x(z, \bar{z})$ furnishes a two-parameter family of metrics labelled by the points $x^\mu \in \mathcal{N}$ and a bundle over $\mathcal{N}$ is obtained whose fibers at each point are isomorphic to $\mathbf{C}^2$ and equipped with a half-flat metric with two Killing vectors.

When $d = 2$ it is possible to construct invariant actions under a subgroup of the $\mathcal{W}$ transformations (symplectic) if, and only if, a $\mathcal{W}$ scalar-density exists, $\tilde{F}(x^\mu, y_\mu)$ so that the action is $\mathcal{W}$ -invariant up to surface terms . In $d > 2$ no such density exists. The reason that the action was solely invariant under a subgroup of $Diff_o(T^* \mathcal{N})$ was due to the fact that a constraint on the gauge parameters $\lambda_{(s)}^{\mu_1 \dots \mu_s}$ had to be imposed as well because under $\mathcal{W}$ transformations given by eqs-(17,19), the action behaves like :

$$\delta S = \int d^2 x \partial_\mu (\Omega^\mu + X) = 0 \Rightarrow X = 0. \quad (24)$$

To *lowest* order in $\Lambda(x^\mu, y_\mu)$ the constraint $X = 0$ reads :

$$\det\left[\frac{\partial^2(\tilde{F} + \Lambda)(x^\mu, y_\mu)}{\partial y_\mu \partial y_\nu}\right] = 1. \quad (25)$$

for $2 + 2$ signature one has -1 on the r.h.s of (25). The above constraint represents *infinitesimal* deformations of the hyper-Kahler geometry with two-Killing vectors, for a deformed Kahler potential $\tilde{F} \rightarrow \tilde{F} + \Lambda$. The constraint $X = 0$ on the gauge parameters is not fully symplectic-diffs invariant like eq- (21) was, it is only invariant under a subgroup of the symplectic-diffs [6]. In the next section we will present a straightforward derivation of all these constraints based on Moyal deformations of SDYM.

Summarizing, invariant actions under a subgroup of the symplectic-diffs can be constructed in terms unconstrained gauge fields, $h_{(s)}^{\mu_1 \dots \mu_s}$, and transformation laws with gauge parameters, $k_{(s)}^{\mu_1 \dots \mu_s}$. Their traces can be removed by means of exploiting the $\mathcal{W}$ -Weyl conformal invariance. These traceless (irreducible) fields and parameters appear in the invariant action and transformation laws *nonlinearly* in the form of $\tilde{h}_{(s)}^{\mu_1 \dots \mu_s}$, and $\lambda_{(s)}^{\mu_1 \dots \mu_s}$. The symmetry algebra of nonchiral W_∞ Gravity is a subalgebra of the $\text{Diff}_o(T^*\mathcal{N})$.

For references on the twistor transform, reductions to W_N Geometry and its connection to Strominger's special geometry [28], finite versus infinitesimal transformations, etc... we refer to Hull's original work [6]. Now we are ready to recast the constraints (21,25) in the language of Fedosov-Moyal quantization.

3.2 W-Geometry from Moyal-Fedosov

From the last two sections we can see that Hull's construction of W_∞ Geometry fits very naturally with the Moyal Self Dual Gravitational equations (14,15) (after a Killing symmetry reduction). The $\hbar = 0$ limit in eqs-(14,15) of the Moyal brackets turns into Poisson brackets and the latter, in turn, can be formulated as a simple determinant :

$$\{\partial_x K_{\tilde{x}, \tilde{y}}(x, y, q, p, \hbar = 0), \partial_y K_{\tilde{x}, \tilde{y}}(x, y, q, p, \hbar = 0)\}_{\text{Poisson}} = 1 \Rightarrow \det\left[\frac{\partial^2 K_{\tilde{x}, \tilde{y}}(x, y, q, p, \hbar = 0)}{\partial \xi^i \partial \bar{\xi}^j}\right] = 1. \quad (26)$$

where $\xi^1; \xi^2$ plus complex conjugates $\bar{\xi}^1; \bar{\xi}^2$ are suitable functions of the x, y, q, p variables. Two Killing symmetry reductions must be subsequently performed in such a way that the x, y, q, p dependence appears solely in the combinations $y^i = \xi^i + \bar{\xi}^i, i = 1, 2$. In this fashion one can make contact with the two-Killing symmetry reductions imposed by Hull [6]. The four variables x, y, q, p admit a natural interpretation in terms of the $z_\mu, \bar{z}_\mu$ variables which described the $\mathbf{C}^2$ fibers of the complexified cotangent space of the two-dim surface, $\mathcal{N}$ at a given point $x^\mu \in \mathcal{N}$. The $\hbar = 0$ limit of the Moyal heavenly equations associated with the two-parameter family of leaves foliating the $6D$ space, after a further

Killing symmetry reduction $y^i = \xi^i + \bar{\xi}^i$, admit a direct correspondence with the constraints (21,23) present in Hull's formulation :

$$K_{\tilde{x},\tilde{y}}(x, y, q, p, \hbar = 0) = K_{\tilde{x},\tilde{y}}(y^i = \xi^i + \bar{\xi}^i, \hbar = 0) \leftrightarrow K_{x^\mu}(z_\mu, \bar{z}_\mu) = K_{x^\mu}(y_\mu = z_\mu + \bar{z}_\mu) = \tilde{F}(x^\mu, y_\mu). \quad (27)$$

Since we were able to identify $K_{\tilde{x},\tilde{y}}(x, y, q, p, \hbar)$ with a two-parameter family of Moyal Kahler potentials , $\Theta(\tilde{x}, \tilde{y}|x, y, q, p, \hbar)$, it follows that Hull's co-metric density function can naturally be embedded into Θ by performing the *double* infinite summation typical of Fedosov's Geometry and recurring to the two Killing symmetry reductions :

$$\Theta(\tilde{x}, \tilde{y}|x, y, q, p, \hbar) = \sum_{2n+l \geq 0} \hbar^n \Theta_{n;i_1, \dots, i_l} (\xi + \bar{\xi})^{i_1} \dots (\xi + \bar{\xi})^{i_l}. \quad (28)$$

the coefficients belong to a one parameter family of smooth tensorial functions of the $\tilde{x}, \tilde{y}$ variables (parametrized by the integer n) :

$$\Theta_{n;i_1, \dots, i_l}(\tilde{x}, \tilde{y}). \quad (29)$$

Hull's co-metric density corresponds solely to the $\hbar^0$ terms :

$$\mathcal{F}(x^\mu, y_\mu) \leftrightarrow \sum_{l \geq 0} (\xi + \bar{\xi})^{i_1} \dots (\xi + \bar{\xi})^{i_l} \Theta_{0;i_1, \dots, i_l}(\tilde{x}, \tilde{y}).. \quad (30)$$

while the higher order $\hbar$ corrections implement in one scoop the Moyal deformation quantization of nonchiral W_∞ gravity. The series in (30) must start with $l = 2$ in order to match the expression for the co-metric density. A truncation to zero of the first two terms in the series is necessary. Futhermore, we have an expansion in contravariant vectors versus Hull's expansion in terms of covariant vectors. Indices are raised and lowered using the two-dim metric. Eq-(30) may be reinterpreted as the zeroth-order terms (in $\hbar$) associated with a Fedosov deformation quantization of the symplectic two-dim manifold with coordinates $\tilde{x}, \tilde{y}$. The tangent vector at the each point should have for coordinates $y^i = \xi^i + \bar{\xi}^i$. This is relevant to the chiral model approach to self dual gravity [10] where instead of starting with a direct Yang-Mills formulation one writes down the Fedosov deformations of WZNW actions.

If the afore-mentioned interpretation is adopted, Hull's action will be the integration of the $\hbar^0$ componet of Θ along a section of the cotangent bundle of the two-dim manifold. Notice that this is *not* the same as taking the Fedosov trace of the zero form, Θ , belonging to the exterior algebra of the Weyl algebra bundle associated to the two-dim symplectic manifold (Riemann surface). Secondly, the W_∞ transformations of the higher spin fields and the matter fields in eqs-(17,19) must not be confused with the Fedosov's noncommutative fibrewise product algebra of the y^i coordinates. Although the W_∞ algebra can be identified with Moyal deformations of the classical area-preserving diffs [24].

Instead of the foliation picture presented above of the $6D$ space into a family of Ricci flat $4D$ leaves, there is yet another interpretation of the relation of the Moyal SDYM equations and Hull's formulation of W_∞ gravity; i.e the constraints (21) on the Kahler

potential. This requires now a dimensional reduction of the original four dimensional spacetime and to extend the construction of **2.2** to the case that the phase space manifold parametrized by q, p is no longer flat. This is attained as follows :

The gauge fields living on the four-dim manifold, R^4 , will now take values in the Weyl algebra bundle, $\mathcal{W}$, constructed over the curved phase space, Γ , whose coordinates are q, p . Now one has $\tilde{A}_\mu(X^\mu|q, p, y^1, y^2; \hbar)$. The zeroth-element is precisely the central section of the algebra $\mathcal{E}(\mathcal{W}_D)$ obtained from the projection : $\sigma(\tilde{A}_\mu) = \tilde{A}_\mu(X^\mu|q, p, 0, 0; \hbar) = A_\mu(X^\mu|q, p; \hbar)$. Fedosov gave the relation which allows one to reconstruct the full $\tilde{A}_\mu$ field from the knowledge of the central section : A_μ . It resembles an expansion in terms of Riemann normal coordinates, see [3,10, 33] for further details. As explained in **2.1** the bijection map, σ yields Moyal star products in the space of central sections inherited from the Fedosov fibrewise products of the flat sections.

Eq-(10) can be obtained from a Lagrangian both in the Moyal and Fedosov case. They represent the deformations of the six dimensional version of the second heavenly equation . The only difference in the Fedosov case is that now one is required to use the definition of the Fedosov trace and to take Fedosov's fibre-wise products instead of Moyal ones. In the flat phase space limit, $\Gamma \rightarrow R^2$, one recovers naturally the Lagrangian of [9] . The trace becomes then an integration w.r.t the q, p coordinates. The original four-dim action will then be an effective $6D$ one : $M^4 \times R^2$ as a result of taking the Fedosov trace :

$$S = \int d^4X \int dqdp \left[-\frac{1}{3} \Theta_o * \{ \partial_x \Theta_o, \partial_y \Theta_o \} + \frac{1}{2} (\partial_x \Theta_o) * (\partial_{\tilde{x}} \Theta_o) + \frac{1}{2} (\partial_y \Theta_o) * (\partial_{\tilde{y}} \Theta_o) \right]. \quad (31)$$

with $x, y, \tilde{x}, \tilde{y}$ being the coordinates of M^4 . The Moyal star product is w.r.t the q, p coordinates of the flat phase space, R^2 . $\Theta_o(X^4|q, p, \hbar)$ is a scalar field living on M^4 and taking values in the space of central sections of the $\mathcal{E}(\mathcal{W}_D)$ algebra. The action yields the equations of motion for the Moyal deformations of the $6D$ version of the second heavenly equation. In the classical $\hbar = 0$ limit, Moyal brackets turn into Poisson ones. A dimensional reduction from $6D$ to $4D$ of the type : $\partial_x = \partial_{\tilde{x}}$ and $\partial_y = \partial_{\tilde{y}}$ furnishes the Park-Hussain second heavenly equations associated with the chiral model approach to Self Dual Gravity. Once again, we see that SDG is an essential geometrical ingredient in these constructions.

Therefore, after an integration of eq-(31) w.r.t the q, p coordinates is performed and a *further* dimensional reduction from M^4 to two-dimensions is taken one can make contact with Hull's action (18). Both views should in principle be equivalent since both lead to Self Dual Gravity. One is expressed in terms of the first heavenly form and the other in terms of the second heavenly one. A Darboux transformation relates the first heavenly equation with the second heavenly one (the chiral models) . Further analysis of these two views : foliations versus reductions and the construction of induced W_∞ gravity from WZNW models [21,43] and its Fedosov deformation quantization will appear elsewhere.

The other constraint (25) that broke the full symplectic-diffs invariance down to a subgroup can also be understood within the framework of Moyal SDYM. The analogy of gauge transformations of the Moyal YM potentials reads :

$$A_M \rightarrow A_M + \partial_M \lambda + \{A_M, \lambda\}_{Moyal}. \quad F_{MN} \rightarrow \{F_{MN}, \lambda\}_{Moyal}. \quad (32a)$$

under gauge transformations the zero curvature (SDYM) conditions are preserved and hence the Moyal Plebanski equations are naturally gauge invariant exactly like it happened to the constraint (21). However, one can see that $\Theta \rightarrow \Theta + \Lambda$ is not always a symmetry of the Moyal SDYM theory. Under shifts in Λ the Moyal YM potentials do *not* transform as (31) but instead :

$$A_{\tilde{x}} = -\partial_y \Theta \Rightarrow A_{\tilde{x}} \rightarrow A_{\tilde{x}} - \partial_y \Lambda. \quad (32b)$$

eq-(32) does not represent a true gauge transformation of the $A_{\tilde{x}}$ Moyal YM potential due to the gauge noncovariant $-\partial_y \Lambda$ piece. One would only have true gauge covariance if, and only if, Λ, λ satisfy the property :

$$-\partial_y \Lambda = -\partial_y \Lambda - \{A_y, \Lambda\}_{Moyal} = \partial_{\tilde{x}} \lambda + \{A_{\tilde{x}}, \lambda\}_{Moyal} = \delta A_{\tilde{x}} \quad (33)$$

which implies the two conditions :

$$\{A_y, \Lambda\}_{Moyal} = 0. \quad -D_y \Lambda = D_{\tilde{x}} \lambda. \quad (34)$$

this is a very restricted condition on the gauge parameter λ and the shift parameter Λ . It is not surprising that the symplectic-diffs invariance is not fully preserved under shifts in Λ of the Moyal Kahler potential Θ . We recall from [9] that A_y could be gauged to zero due to the zero curvature condition that furnishes the Moyal SDYM equations in R^4 . Hence, the gauge $A_y = 0$ obeys one of the conditions. However, there still remains the other condition restricting the parameters λ, Λ in a highly nonlinear manner in terms of the remaining Moyal YM potentials. Similar considerations apply for the other $A_{\tilde{y}}$ Moyal YM potential.

IV. Conclusion

We have shown that Hull's formulation of nonchiral W_∞ Geometry fits in very naturally inside a larger picture : Moyal-Fedosov Geometry. Hull's constraints (21,25) have a natural interpretation in the Fedosov-Moyal quantization program. Furthermore, the inclusion of all the powers in $\hbar$ implements in a straightforward fashion the deformation quantization of nonchiral W_∞ gravity. In general one must consider a Yang-Mills like formulation of $\mathcal{W}$ -geometry based on a $\mathcal{W}$ -valued connection one-form belonging to the exterior algebra of the Weyl-algebra bundle associated with the symplectic manifold. Some of the advantages of this formulation of $\mathcal{W}$ -geometry are the following ,

- The incorporation of many bosonic fields $\phi^i, i = 1, 2, 3, \dots, D$ is straightforward in Fedosov's Geometry : the coefficients in the expansion (1) are matrix valued; i.e. the coefficients take values in the bundle $\text{Hom}(E, E)$ where E is a vector bundle over $\mathcal{M}$. For more details on this see [3]. The scalars ϕ^i represent the embedding coordinates of the world sheet in a target spacetime background of dimension D . Since these fields are reinterpreted as matrix valued sections with a noncommutative fibrewise Fedosov product it is clear why the embedding coordinates of the string world sheet inherit a noncommutative product structure !

- Quantization of p -branes [37] should be achieved using the Zariski product [34] where deformation quantization methods for higher dimensional generalizations of symplectic geometry are achieved : the so-called Nambu-Poisson Hamiltonian Mechanics. Deformation quantization of Poisson manifolds has also been discussed by [35,36].

- We hope that the Moyal-Fedosov deformation quantization approach to $\mathcal{W}$ Geometry will provide many new insights into the nonperturbative structure of string theory. In particular the role of W_∞ strings [8]. Paraphrasing Fairlie [16] : Moyal stands for “M”; to-day we advocate that M-theory “stands upside down” for $\mathcal{W}$. Important objections why W geometry is still far from being understood have been recently raised by [39].

- Zucchini has shown that $4D$ Conformal Field Theory can naturally be formulated in real four-folds (Kulkarni four-folds) endowed with an integrable quaternionic structure [26] and a $4D$ extension of $2D$ CFT on Riemann surfaces was constructed. Quaternionic (Fueter) analyticity played the role of $2D$ holomorphic analyticity. It is warranted to explore further the connections between the self duality and $\mathcal{W}$ -Weyl conformal invariance properties of W_∞ Geometry and Zucchini’s Quaternionic Geometry of $4D$ Conformal Field Theory. $4D$ generalizations of the $2D$ WZNW models were studied by [32]. Earlier work in that direction was provided by Park [22]. The fact that quaternions are noncommutative points in the right direction. The non-associative character of octonions suggests that $8D$ Non Associative Geometries are no longer speculative figments of the imagination but should also come to play an important role in physics.

- The view advocated here of $\mathcal{W}$ geometry as flat foliations in higher dimensions may have an important relation with Zois [38] proposal for a nonperturbative Lagrangian of M theory in $11D$ in terms of characteristic classes of flat foliations (although in odd dimensions) .

- A forthcoming project involves to write down Yang-Mills types of action characterizing the higher spin field dynamics. In particular to establish the connection to Vasiliev’s work [40] : Higher-spin gauge theories in four, three and two-dimensions and interactions of matter fields based on deformed oscillator algebras have been studied by Vasiliev and others . A reformulation of the dynamical equations of motion, called “the unfolded formulation “, in a form of a zero curvature condition and a covariant-constancy condition imposed on an infinite collection of zero-forms allowed Vasiliev to describe all spacetime derivatives of all the dynamical fields and reconstruct these fields by analyticity in some neighbourhood of a fixed point. There are many similarities with the work of [40] and ours : a zero curvature condition is imposed; a star product also appears in order to describe the nonlinear dynamics; a deformed oscillator algebra realizing the universal enveloping algebra of symplectic groups is essential; What is required then is to integrate Vasiliev’s formulation in the Moyal-Fedosov program.

Acknowledgments

We are indebted to George Chapline for many discussions .

REFERENCES

1. F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz and D. Sternheimer , “ Deformation Theory and Quantization “ Ann. Phys. **111** (1978) 61.
2. J. Moyal, “Quantum Mechanics as a Statistical Theory”.

- Proc. Cam. Phil. Soc. **45** (1945) 99
- E. Wigner, “ Quantum corrections for thermodynamic equilibrium “
Phys. Rev. **40** (1932) 749.
- H. Groenewold, “ On the principles of elementary quantum mechanics
“ Physica **12** (1946) 405.
- H. Weyl, “ Quantum mechanics and the theory of groups” Z. Phys. **46** (1927) 1.
3. B. Fedosov, “ A simple geometrical construction of deformation quantization”
J. Diff. Geometry, **40** (1994) 213.
4. T. Deroli, A. Verciu, “ W_∞ -covariance of the
Weyl-Wigner-Groenewold-Moyal quantization “ J. Math. Phys. **38** (11) (1997) 5515.
5. P. Bouwknegt, K. Schouetens, “ W-symmetry in Conformal Field Theory”.
Phys. Reports **223** (1993) 183-276. hep-th/9210010.
6. C. Hull, “ The Geometry of W-gravity” Phys. Lett **B 269** (1991) 257.
“ $\mathcal{W}$ -Geometry “ , hep-th/9211113.
“Geometry and $\mathcal{W}$ -gravity “ hep-th/9301074.
7. C. Castro, “ A Moyal quantization of the continous Toda field”
Phys. Lett **B 413** (1997) 53.
“ $SU(\infty)$ (super) gauge theories and self dual (super) gravity” .
J. Math. Phys **34** (1993) 681.
“ On W gravity, $N = 2$ strings and $2 + 2$ $SU(\infty)$ Yang-Mills instantons” .
J. Math. Phys **35** (1994) 3013
8. C. Castro, “ $D = 11$ supermembrane instantons, W_∞ strings and the
super Toda molecule”. J. of Chaos, Solitons and Fractals, **7** (7) (1996) 711,
“ The non-critical W_∞ string sector of the self dual membrane” hep-th/9612160.
9. J. Plebanski, M. Przanowski, “ The Lagrangian of Self
Dual Gravity as the limit of the SDYM Lagrangian “ . Phys. Lett **A 212** (1996) 22
10. H. Garcia-Compean, J. Plebanski and M. Przanowski : “ Geometry associated
with SDYM and the chiral model approach to Self Dual Gravity”. hep-th/9702046.
11. A. Connes, M. Douglas, A. Schwarz, “Non Commutative Geometry and Matrix
Theory : Compactification on Tori” hep-th/9711162.
12. P. Ming Ho, Y.S. Wu, “ Non Commutative Gauge Theories in Matrix Models “ .
hep-th/9801147.
13. F. Lizzi, R. Szabo : “ Non Commutative Geometry and Spacetime Gauge
Symmetries of String Theory “ hep-th/9712206.
14. J. Froehlich, O. Grandjean, A. Recknagel : “ Supersymmetric Quantum Theory,
Non Commutative Geometry and Gravitation ”. hep-th/9706132.
15. T. Banks, W. Fischler, S. Shenker, L. Susskind, “ M theory as a Matrix
theory : a conjecture “ Phys. Rev. **D 55** (1997) 112.
16. D. Fairlie, “ Moyal Brackets in M Theory”. hep-th/9707190.
17. E. Gozzi, M. Reuter , “ Quantum Deformed Geometry in Phase Space”
DESY-92-191 .
18. I. Strachan, “ The Moyal algebra and integrable
deformations of the self dual Einstein equations “ . Phys. Lett. **B 283** (1992) 63.
19. K. Takasaki, “ Dressing operator to Moyal algebraic formulation of

- Self Dual gravity” J. Geom. and Phys. **14** (1994) 111.
20. H. Ooguri, C. Vafa,
 “ Geometry of $N = 2$ strings” Nuc. Phys **B 361** (1991) 469.
21. G. Chapline, K. Yagamishi,
 “ Induced $4D$ self dual quantum gravity : a W_∞ algebraic approach”.
 Class. Quantum. Grav. **8** (1991) 427.
- “ A Hamiltonian structure of the KP hierarchy , $W_{1+\infty}$ algebra and
 self dual gravity.” Phys. Lett. **B 259** (1991) 436.
22. Q.H. Park, “ Extended conformal symmetry in real heavens “
 Phys. Lett. **B 236** (1990) 429.
23. I. Bakas, “ The large N limit of extended conformal symmetries “
 Phys. Lett. **B 228** (1989) 57.
24. D. Fairlie, J. Nuyts, “ Deformations and renormalizations of W_∞
 algebras “ Comm. Math. Phys. **134** (1990) 413.
- I. Bakas, E. Kiritsis, B. Khesin, “ The Logarithm Operator in Conformal
 Field Theory and W_∞ algebras “ Comm. Math. Phys. **151** (1993) 233
25. A. Jevicki , “ Matrix Models, Open Strings and Quantization of
 Membranes “ hep-th/9607187.
26. R. Zucchini, “ The Quaternionic Geometry of $4D$ Conformal Field Theory “
 gr-qc/9707048.
27. C. Castro, J. Plebanski : “ The Generalized Moyal Nahm and
 Continous Moyal Toda Equations” hep-th/9710041.
28. A. Strominger , “ Special Geometry “ Comm. Math. Phys **133** (1990) 163.
29. E. Martinec, “ M Theory and $N=2$ Strings “ hep-th/9710122.
30. C. Devchand, O. Lechtenfeld : “ On The Self Dual Geometry of $N=2$ Strings.
 hep-th/9801117.
31. S. Ketov, “ From $N=2$ Strings to F, M Theories “ hep-th/9606142.
32. A. Lesov, G. Moore, N. Nekrasov and S. Shatashvili, :
 “ Four Dimensional Avatars of two-dim RCFT “ hep-th/9509151.
33. M. Bordemann , N. Neumeier, S. Waldmann : “ Homogeneous Fedosov Star
 Products on Cotangent Bundles I : “ Weyl Ordering with Differential Operators
 Representation “ q-alg/9707030.
- “ Homogeneous Fedosov Star Products on Cotangent Bundles II :
 GNS Representations, the WKB expansion and Applications “ q-alg/9711016.
- P. Xu : “ Fedosov star products and quantum momentum maps “ q-alg/9608006.
- A. Astashkevich, “ On Fedosov’s quantization of semisimple coadjoint orbits” MIT
 Ph.D Thesis. Math. Dept. 1996. ‘
- O. Kravchenko : “ Deformation Quantization of Symplectic Fibrations”.
 math.QA/9802070. “
34. G. Dito, M. Flato, D. Sternheimer and L. Takhtajan :
 “ The deformation quantization and Nambu Mechanics “ hep-th/9602016.
35. M. Kontsevich, “ Deformation Quantization of Poisson Manifolds I “
 q-alg/9709040.
36. A. Voronov, “ Quantizing Poisson Manifolds “ q-alg/9701017.

37. C. Castro, “ p-branes as Composite Antisymmetric Tensor Field Theories “
Int. Jour. Mod. Phys. **A 13** (8) (1998) 1263-1292.
38. I. Zois : “ On search for the M-theory Lagrangian “ hep-th/9703067.
39. S. Lazzarini, “ Flat Complex Vector Bundles , the Beltrami Differential and
W- Algebras “ hep-th/9802083.
40. M. Vasiliev : “ Deformed Oscillator algebras and higher spin gauge
interactions of matter fields in $2 + 1$ dimensions “ hep-th/9712246.
“ Higher-spin gauge theories in four, three, and two dimensions “
hep-th/9611024.
41. T. Ivanova , A. Popov, “Some new integrable equations from the Self Dual
Yang-Mills equations “ Phys. Lett **A 205** (1995) 158. “
A. Popov , “Self Dual Yang-Mills symmetries and moduli spaces
“ hep-th/9803183.
T. Ivanova , “ On infinitesimal symmetries of the self dual Yang-Mills
equations “ physics/9803028.
42. M. Saveliev, A. Vershik : “ Continual analogs of contragredient Lie
Algebras “ Comm. Math. Phys. **126** (1989) 367.
43. E. Nissimov, S. Pacheva, I. Vaysburd : “ Induced W_∞ Gravity from a
WZNW Model “ RI-92 (Racach Institute) and BGU-92 (Ben Gurion University)
preprints.